



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44	A1	(11) International Publication Number: WO 00/27397 (43) International Publication Date: 18 May 2000 (18.05.00)
(21) International Application Number: PCT/GB98/03336 (22) International Filing Date: 6 November 1998 (06.11.98) (71) Applicants: GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; D-55216 Ingelheim am Rhein (DE). (72) Inventor: GAVIRAGHI, Giovanni; Glaxo Wellcome SpA, Via A Fleming, 2, I-37135 Verona (IT). (74) Agent: FILLER, Wendy, A.; Glaxo Wellcome plc, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ANTIHYPERTENSIVE MEDICAMENTS CONTAINING LACIDIPINE AND TELMISARTAN (57) Abstract Combinations comprising diethyl (E) -4-[2-[(tert-butylcarbonyl)vinyl]phenyl-1,4-dihydro-2,6-dimethylpyridine-3,5 dicarboxylate(lacidipine) and 4'-[[2-n-propyl-4-methyl-6-(1-methylbenzimidazol-2-yl)-benzimidazol-1-yl]-methyl]-biphenyl-2-carboxylic acid (telmisartan), pharmaceutical compositions containing said combinations and their use in the treatment of cardiovascular disorders including hypertension.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ANTIHYPERTENSIVE MEDICAMENTS CONTAINING LACIDIPINE AND TELMISARTAN

The present invention relates to therapeutic combinations comprising diethyl (E)-4-[2-[(tert-butylcarbonyl)vinyl]phenyl-1,4-dihydro-2,6-dimethylpyridine-3,5-dicarboxylate(lacidipine) and 4'-[[2-n-propyl-4-methyl-6-(1-methylbenzimidazol-2-yl)-benzimidazol-1-yl]-methyl]-biphenyl-2-carboxylic acid (telmisartan), to pharmaceutical compositions containing said combinations and their use in the treatment of cardiovascular disorders including hypertension.

Lacidipine, which is described in British patent no. 2164336, is a potent long acting calcium antagonist which is particularly useful for treating hypertension. The compound may be also useful for the treatment of other cardiovascular disorders including atherosclerosis, peripheral vascular disease, ischaemic heart disease and congestive heart failure.

Telmisartan, which is described in European patent no. 0502314, is an angiotensin-II-antagonist which is useful for treating hypertension and cardiac insufficiency and for treating other cardiovascular disorders including ischaemic peripheral circulation disorders, myocardial ischaemia (angina).

European patent no. 0502314 teaches that the angiotensin-II-antagonists described therein may be administered in combination with other active substances including calcium antagonists. There is however no specific disclosure of such combinations with lacidipine.

We have found that the combination of lacidipine and telmisartan provides a useful and unexpectedly advantageous combination for the treatment of cardiovascular disorders, such as hypertension, atherosclerosis and ischaemic heart disease.

In particular it has now been found that by combining lacidipine and telmisartan, a synergistic antihypertensive effect is achieved.

It is a feature of this invention that the use of such a drug combination will provide one or more of the following effects: synergistic antihypertensive effects, antihypertensive effect over a longer period and/or allow a better management of any potential drug-related side effects.

Furthermore, the improvement of blood pressure control achieved by using such a drug combination may afford a better protection from the associated diseases which are induced by hypertension.

According to one aspect of the invention there is provided a combination comprising lacidipine and telmisartan or a physiologically functional derivative thereof and more particularly a combination comprising lacidipine and telmisartan.

As used herein, the term "*physiologically functional derivative*" includes any physiologically acceptable solvate, salt, ester, salt of such ester, or solvates of any such salt or ester, of telmisartan.

Preferred esters in accordance with the invention are independently selected from the following group: (1) carboxylic acid esters in which the non-carbonyl moiety of the carboxylic acid portion of the ester grouping is selected from straight or branched chain alkyl (for example, methyl, *n*-propyl, *t*-butyl, or *n*-butyl), cycloalkyl, alkoxyalkyl (for example, methoxymethyl), aralkyl (for example, benzyl), aryloxyalkyl (for example, phenoxymethyl), aryl (for example, phenyl optionally substituted by, for example, halogen, C₁₋₄ alkyl, or C₁₋₄ alkoxy), or amino; (2) sulphonate esters, such as alkyl- or aralkylsulphonyl (for example, methanesulphonyl); (3) amino acid esters (for example, L-valyl or L-isoleucyl); and (4) phosphonate esters. In such esters, unless otherwise specified, any

alkyl moiety present advantageously contains from 1 to 18 carbon atoms, particularly from 1 to 6 carbon atoms, more particularly from 1 to 4 carbon atoms. Any cycloalkyl moiety present in such esters advantageously contains from 3 to 6 carbon atoms. Any aryl moiety present in such esters
5 advantageously comprises a phenyl group. Any reference to any of the above compounds also includes a reference to a physiologically acceptable salt thereof.

Examples of physiologically acceptable salts include salts derived from an
10 appropriate base, such as an alkali metal (for example, sodium), an alkaline earth (for example, magnesium), ammonium and NX_4^+ (wherein X is C_{1-4} alkyl) or

ammonium salts, formed with amino acids (e.g lysine and arginine) and organic bases (e.g procaine, phenylbenzylamine, ethanolamine and N-methyl
15 glucosamine).

Salts of acids or bases which are not physiologically acceptable may also find use, for example, in the preparation or purification of a physiologically acceptable compound. All salts, whether or not derived from a physiologically acceptable acid or base, are within the scope of the present invention.
20

The present invention thus provides a method for the treatment of hypertension in a mammal including a human, which comprises treating said animal with a therapeutically effective amount of a combination of lacidipine and telmisartan or a physiologically functional derivative thereof.
25

Reference herein to treatment extends to prophylaxis as well as the treatment of established hypertension or symptoms.

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations or sequentially. If there is sequential administration, the delay in administering the second and any subsequent active ingredient should not be such as to lose the benefit of a synergistic therapeutic effect of the combination of the active ingredients. It will also be understood that the compounds of the combination or the physiologically functional derivatives of any thereof, whether presented simultaneously or sequentially, may be administered individually or in multiples or in any combination thereof.

According to another aspect, the present invention provides the use of lacidipine in the manufacture of a medicament for administration simultaneously or sequentially with telmisartan or a physiologically functional derivative thereof for the treatment and/or prophylaxis of hypertension.

The synergistic effects of the combination of lacidipine and telmisartan may be seen over a wide ratio of combinations, for example, of 1: 100 to 1: 1, such as 1:50 to 1:2 (by weight), preferably of 1:40 to 1:3.33 (by weight). Examples of such combinations include those wherein the ratio (by weight) of lacidipine to telmisartan is 1:5; 1:10, 1:20; 1:40; 1:6.67 or 1:13.33. Conveniently each compound will be employed in the combination in an amount at which it exhibits an antihypertensive effect when used alone.

The amount of a combination of lacidipine and telmisartan required to be effective as antihypertensive may, of course, vary and is ultimately at the discretion of the medical practitioner. The factors to be considered include the route of administration and nature of the formulation, the animal's body weight, age and general condition and the nature and severity of the disease to be treated.

In general a suitable dose of lacidipine for administration to a human for the treatment of hypertension may be in the range of 0.1 to 10 mg per day, preferably in the range of 1 to 6 mg per day and most preferably in the range 2-
5 6 mg per day. Lacidipine is advantageously administered by oral route once a day.

In general, a suitable dose of telmisartan for administration to a human may be in the range of 5 to 120 mg per day, advantageously in the range of 20 to 80
10 mg per day. Telmisartan is advantageously administered by oral route once a day.

Unless otherwise indicated all weights of active ingredients are calculated in terms of the drug *per se*. The desired dose may preferably be presented as one,
15 two, three, four, five, six or more sub-doses administered at appropriate intervals throughout the day. Conveniently lacidipine and telmisartan are administered as a single daily dose.

20 The components of the combination which may be referred to as active ingredients may be administered for therapy to an animal e.g. a mammal including a human in a conventional manner.

While it is possible for the active ingredients of the combination to be
25 administered as the raw chemical it is preferable to present them as a pharmaceutical formulation. Pharmaceutical formulations according to the present invention comprise a combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. The carrier(s) must be acceptable in the

sense of being compatible with the other ingredients of the formula and not deleterious to the recipient thereof. When the individual components of the combination are administered separately they are generally each presented as a pharmaceutical formulation. The references hereinafter to formulations refer,
5 unless otherwise stated, to formulations containing either the combination or a component thereof.

A combination of lacidipine and telmisartan or a physiologically functional derivative thereof may conveniently be presented as a pharmaceutical
10 formulation in a unitary dosage form. A convenient unitary dosage formulation contains lacidipine in an amount from 1mg to 6 mg and telmisartan in an amount from 10 mg to 100 mg.

A particularly convenient unitary dosage formulation contains lacidipine in an amount from 2 mg to 6 mg, more particularly in an amount from 2mg to 4 mg,
15 and telmisartan in amount from 20mg to 80 mg.

Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package, usually a blister pack. Patient packs have an advantage over traditional prescriptions,
20 where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions and, therefore, lead generally to more successful
25 treatment.

It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each formulation,

containing within a package insert instructing the patient to the correct use of the invention is a desirable additional feature of this invention.

5 According to a further aspect of the invention provided is a multiple, for example, double or triple, pack comprising at least lacidipine and telmisartan or a physiologically functional derivative thereof and an information insert containing directions on the use of the combination of the invention.

10 Formulations include those suitable for oral, rectal, nasal, topical (including transdermal, buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods represent a further feature of the present invention and include the step of
15 bringing into association the active ingredients with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.

20 Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, caplets, cachets or tablets each containing a predetermined amount of the active ingredients; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or
25 as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in

a suitable machine the active ingredients in a free-flowing form such as a powder or granules, optionally mixed with a binder (e.g. povidone, gelatin, hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (e.g. sodium starch glycollate, sodium croscarmellose cross-linked povidone, cross-linked sodium carboxymethyl cellulose) surface-active or dispersing agent. Molded tablets may be made by molding a mixture of the powdered compound moistened with an inert liquid diluent in a suitable machine. The tablets may optionally be coated or scored any may be formulated so as to provide slow or controlled release of the active ingredients therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredients in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier. Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or polyethylene glycols.

Topical administration may also be by means of a transdermal iontophoretic device.

Formulations suitable for vaginal administration may be presented as tablets, pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Pharmaceutical formulations suitable for rectal administration wherein the carrier is a solid are most preferably presented as unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by admixture of the active combination with the softened or melted carrier(s) followed by chilling and shaping in molds.

Formulations suitable for parenteral administration include aqueous and nonaqueous isotonic sterile injection solutions which may contain anti-oxidants, buffers, preservatives and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents; and liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example, those suitable for oral administration may include such further agents as sweeteners, thickeners and flavoring agents.

The pharmaceutical composition of the invention containing the two active ingredients may be prepared according to conventional techniques well known

in the pharmaceutical industry. Thus, for example the lacidipine and telmisartan may be admixed together with suitable excipients such as those described above for the formulation of each of the active ingredients separately. Tablets may be prepared, for example by direct compression of such a mixture or using
5 other conventional methods. Bilayer tablets may be prepared according to conventional procedure. Thus, for example, by separately compressing the two blends in a suitable tableting machine with two filling stations. Capsules may be prepared by filling the blend along with suitable excipients into gelatin capsules, using a suitable filling machine. Controlled release forms for oral or rectal
10 administration may be formulated in a conventional manner associated with controlled release forms.

Biological data:

The advantageous profile of antihypertensive activity obtained by the
15 administration of lacidipine with telmisartan may be demonstrated in the male spontaneously hypertensive rats.

In this test lacidipine (0.2mg/kg), telmisartan (1mg/kg) and a combination of lacidipine (0.2mg/kg) and telmisartan (1mg/kg) were administered as a
20 suspension in 0.5% MethocelTM (Hydroxypropyl methyl cellulose) by oral gavage to male spontaneously hypertensive rats.

After dosing, mean blood pressure (MBP) and heart rate (HR) variations were calculated at fixed intervals and expressed as a percentage of pre-drug values.
25 The results obtained three hours after administration of lacidipine and telmisartan, alone or in combination, are summarised below:

Compound			
	lacidipine	telmisartan	lacidipine

	0.2mg/kg	1mg/kg	0.2mg/kg telmisartan 1mg/kg
MBP	-13.8 ± 5.4	-4.5 ± 8.2	-30.5 ± 5.0
HR	1.9 ± 6.4	2.7 ± 17.0	12.6 ± 23.1

5 The reduction in mean blood pressure with the combination of lacidipine and telmisartan was significantly greater than was to be expected and this was also achieved without a significant effect on the heart rate. Duration of action studies also established that the antihypertensive effect of the lacidipine-telmisartan combination was clearly longer lasting than the effect induced by lacidipine and telmisartan alone.

10 The compounds of the combination of the present invention may be obtained in a conventional manner.

Lacidipine may be prepared by the method described in British Patent N° 2164336 which is incorporated herein by reference hereto.

15 Telmisartan or a physiologically functional derivative thereof may be prepared by the method described in European Patent N°502314 which is incorporated herein by reference or by known methods described for analogous compounds

20 For co-administration the lacidipine and telmisartan may be formulated in a conventional manner. Thus for example lacidipine may be formulated as described in British Patent N° 2164336 and telmisartan may be formulated as described in European Patent N° 0502314.

25 In a preferred aspect of the invention lacidipine and telmisartan are formulated in a single pharmaceutical composition.

In order that this aspect of the invention may be more fully understood the following examples are given by way of illustration only.

5 **TABLET FORMULATIONS:**

Example 1

The following formulation was prepared by mixing lacidipine granulated containing monohydrate lactose and telmisartan spray dried granulate with
10 sorbitol, followed by addition of magnesium stearate and compression.

	mg/tablet
Lacidipine	4
Telmisartan	40
Monohydrate Lactose	197
Sodium Hydroxide	3.36
Meglumine	12
Povidone	52
Sorbitol	184
Magnesium Stearate	7.5

15 **Example 2**

The following formulation was prepared by mixing telmisartan spray dried granule with sorbitol and magnesium stearate. Then lacidipine granule was mixed with the remaining magnesium stearate and eventually with sorbitol. The

two blends were separately compressed in a suitable tableting machine with two filling stations to produce bilayer tablets.

5

	mg/tablet
Telmisartan	40
Lacidipine	4
Povidone	40
Monohydrate Lactose	197
Sodium Hydroxide	3.36
Meglumine	12
Povidone	12
Sorbitol	184.1

10

The following formulations (Examples 3a-3d) may be prepared by mixing a granulate containing lacidipine, sorbitol and povidone with telmisartan spray dried granulate, sorbitol, followed by addition of magnesium stearate and compression.

15

Example 3a

	mg/tablet
Telmisartan	40
Lacidipine	4
Povidone	52

14

Sorbitol	116
Sodium Hydroxide	3.36
Meglumine	12
Sorbitol	117.64
Magnesium Stearate	5

Example 3b

	mg/tablet
Telmisartan	20
Lacidipine	2
Povidone	26
Sorbitol	138
Sodium Hydroxide	1.68
Meglumine	6
Sorbitol	151.32
Magnesium Stearate	5

5

Example 3c

10	mg/tablet
Telmisartan	80
Lacidipine	2
Povidone	44
Sorbitol	138

15

Sodium Hydroxide	6.72
Meglumine	24
Sorbitol	50.28
Magnesium Stearate	5

Example 3d

	mg/tablet
Telmisartan	80
Lacidipine	6
Povidone	84
Sorbitol	94
Sodium Hydroxide	6.72
Meglumine	24
Sorbitol	50.28
Magnesium Stearate	5

5

Example 4

The following formulation was prepared by granulating telmisartan spray dried granule and sorbitol with lacidipine and povidone followed by addition of sorbitol and 10 magnesium stearate and compression.

	mg/tablet
Lacidipine	4
Povidone	40

16

Sorbitol	88.64
Telmisartan	40
Sodium Hydroxide	3.36
Meglumine	12
Povidone	12
Sorbitol	295
Magnesium Stearate	5

CLAIMS

1. A combination comprising diethyl (E) -4-[2-[(tert-butylcarbonyl)vinyl]phenyl-
1,4-dihydro-2,6-dimethylpyridine-3,5 dicarboxylate (lacidipine) and 4'-[[2-n-
5 propyl-4-methyl-6-(1-methylbenzimidazol-2-yl)-benzimidazol-1-yl]-methyl]-
biphenyl-2-carboxylic acid (telmisartan) or a physiologically functional
derivative thereof.
2. A combination according to claim 1 for use in medical therapy.
- 10 3. A combination according to claim 1 for use in the treatment and or
prophylaxis of hypertension.
4. A combination according to claim 1 wherein the ratio of lacidipine to
15 telmisartan is from 1:100 to 1:1 by weight.
5. A method for the treatment of hypertension in a mammal including a human,
which comprises treating said animal with a therapeutically effective amount
of a combination of as claimed in claim 1.
- 20 6. A method according to claim 5 wherein the combination is administered as a
single combined formulation.
7. A pharmaceutical formulation comprising a combination according to claim 1
25 together with one or more pharmaceutically acceptable carriers or excipients.
8. A pharmaceutical formulation according to claim 7 in unit dosage form.

9. The use of lacidipine in the manufacture of a medicament for administration simultaneously or sequentially with telmisartan or a physiologically functionally derivative thereof for the treatment and/or prophylaxis of hypertension.

5

10. A patient pack comprising lacidipine and telmisartan or a physiologically functional derivative thereof.

10

15

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/GB 98/03336

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 502 314 A (THOMAE GMBH DR K) 9 September 1992 cited in the application see page 20, line 46 - page 21, line 6; claims 1,5,8 ---	1-3,5-10
Y	WO 96 07400 A (ASTRA AB) 14 March 1996 see page 6, line 4-22 see page 7, line 4-5; claims 1,5-13,21-27 ---	1-10
Y	WO 96 19233 A (OMEROS MED SYS INC) 27 June 1996 see page 9, line 7-25; examples 4-6 see page 10, line 11-20 see page 28, line 5 - page 31, line 4 --- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 July 1999

Date of mailing of the international search report

12/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kanbier, D

INTERNATIONAL SEARCH REPORT

Inter nal Application No

PCT/GB 98/03336

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 36874 A (SMITHKLINE BEECHAM CORP) 9 October 1997 see page 7, line 30 - page 8, line 9; claims 11,16,17 -----	1-10
A	WO 98 40067 A (SANOFI SA) 17 September 1998 see page 10, line 11 - page 11, line 22 -----	1-10
A	WO 98 30216 A (MERCK & CO INC) 16 July 1998 see claims -----	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 98/03336

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 1-10
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 98/03336

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0502314	A	09-09-1992	DE 4103492 A	08-10-1992
			DE 4117121 A	17-12-1992
			DE 4137812 A	19-05-1993
			AT 166346 T	15-06-1998
			AU 655794 B	12-01-1995
			AU 1070792 A	13-08-1992
			CA 2060624 A	07-08-1992
			CS 9200306 A	12-08-1993
			DE 59209330 D	25-06-1998
			ES 2118095 T	16-09-1998
			FI 920486 A	07-08-1992
			HR 940752 A	30-04-1997
			HU 9500157 A	28-11-1995
			IL 100864 A	19-01-1996
			JP 2709225 B	04-02-1998
			JP 4346978 A	02-12-1992
			MX 9200509 A	01-08-1992
			NO 301585 B	17-11-1997
			NZ 241515 A	26-10-1994
			PL 169675 B	30-08-1996
			SG 50481 A	20-07-1998
			SI 9210098 A	31-12-1994
			SK 279261 B	05-08-1998
			RU 2053229 C	27-01-1996
			US 5594003 A	14-01-1997
			US 5602127 A	11-02-1997
			US 5614519 A	25-03-1997
			US 5591762 A	07-01-1997
			ZA 9200816 A	05-07-1993
			EP 0543263 A	26-05-1993
WO 9607400	A	14-03-1996	AU 695865 B	27-08-1998
			AU 3402995 A	27-03-1996
			CA 2174338 A	14-03-1996
			CN 1134665 A	30-10-1996
			CZ 9601173 A	11-09-1996
			DE 726766 T	30-01-1997
			EP 0726766 A	21-08-1996
			ES 2091173 T	01-11-1996
			FI 961863 A	02-05-1996
			GR 96300055 T	31-10-1996
			HU 76049 A	30-06-1997
			JP 9504806 T	13-05-1997
			NO 961726 A	29-04-1996
			NZ 292298 A	26-02-1998
			PL 314084 A	19-08-1996
			SK 55896 A	05-03-1997
			TR 960186 A	21-06-1996
			ZA 9507043 A	04-03-1996
WO 9619233	A	27-06-1996	AU 4467396 A	10-07-1996
			BR 9509985 A	03-11-1998
			CA 2206119 A	27-06-1996
			CN 1175213 A	04-03-1998
			EP 0799051 A	08-10-1997
			JP 10510540 T	13-10-1998
			NO 972687 A	07-08-1997
			US 5820583 A	13-10-1998

INTERNATIONAL SEARCH REPORT

information on patent family members

Int. Application No

PCT/GB 98/03336

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9619233 A		US 5800385 A	01-09-1998
		US 5860950 A	19-01-1999
		US 5858017 A	12-01-1999
WO 9736874 A	09-10-1997	AU 2547097 A	22-10-1997
		CZ 9803101 A	17-03-1999
		EP 0889880 A	13-01-1999
		NO 984503 A	28-09-1998
		NZ 332008 A	28-05-1999
		PL 329046 A	01-03-1999
WO 9840067 A	17-09-1998	FR 2760364 A	11-09-1998
		AU 6840098 A	29-09-1998
WO 9830216 A	16-07-1998	AU 6021898 A	03-08-1998
		ZA 9800163 A	10-07-1998